

Endo logo (scripted)


ENDO LABORATORIES

**SYMMETREL®**  
(Amantadine Hydrochloride)  
Tablets and Syrup, USP

**R<sub>x</sub> only**

**DESCRIPTION**

SYMMETREL is designated generically as amantadine hydrochloride and chemically as 1-adamantanamine hydrochloride.



Amantadine hydrochloride is a stable white or nearly white crystalline powder, freely soluble in water and soluble in alcohol and in chloroform.

Amantadine hydrochloride has pharmacological actions as both an anti-Parkinson and an antiviral drug.

SYMMETREL is available in tablets and syrup.

Each tablet intended for oral administration contains 100 mg amantadine hydrochloride and has the following inactive ingredients: hydroxypropyl methylcellulose, magnesium stearate, microcrystalline cellulose, sodium starch glycolate, FD&C Yellow No. 6.

SYMMETREL syrup contains 50 mg of amantadine hydrochloride per 5 mL and has the following inactive ingredients: artificial raspberry flavor, citric acid, methylparaben, propylparaben, and sorbitol solution.

**CLINICAL PHARMACOLOGY**

**Pharmacodynamics**

**Mechanism of Action: Antiviral** The mechanism by which amantadine exerts its antiviral activity is not clearly understood. It appears to mainly prevent the release of infectious viral nucleic acid into the host cell by interfering with the function of the transmembrane domain of the viral M2 protein. In certain cases, amantadine is also known to prevent virus assembly during virus replication. It does not appear to interfere with the immunogenicity of inactivated influenza A virus vaccine.

**Antiviral Activity:** Amantadine inhibits the replication of influenza A virus isolates from each of the subtypes, i.e., H1N1, H2N2 and H3N2. It has very little or no activity against influenza B virus

isolates. A quantitative relationship between the *in vitro* susceptibility of influenza A virus to amantadine and the clinical response to therapy has not been established in man. Sensitivity test results, expressed as the concentration of amantadine required to inhibit by 50% the growth of virus (ED<sub>50</sub>) in tissue culture vary greatly (from 0.1 µg/mL to 25.0 µg/mL) depending upon the assay protocol used, size of virus inoculum, isolates of influenza A virus strains tested, and the cell type used. Host cells in tissue culture readily tolerated amantadine up to a concentration of 100 µg/mL.

**Drug Resistance:** Influenza A variants with reduced *in vitro* sensitivity to amantadine have been isolated from epidemic strains in areas where adamantane derivatives are being used. Influenza viruses with reduced *in vitro* sensitivity have been shown to be transmissible and to cause typical influenza illness. The quantitative relationship between the *in vitro* sensitivity of influenza A variants to amantadine and the clinical response to therapy has not been established.

**Mechanism of Action: Parkinson's Disease** The mechanism of action of amantadine in the treatment of Parkinson's disease and drug-induced extrapyramidal reactions is not known. Data from animal studies have either shown or suggested SYMMETREL:

- (a) To enhance extracellular concentrations of dopamine by increasing dopamine release or decreasing reuptake of dopamine into presynaptic neurons;
- (b) To stimulate the dopamine receptor itself or drive the post synaptic dopaminergic system to a more dopamine sensitive status.

However, doses employed in the animal studies were often of a magnitude greater than the clinically therapeutic doses. More recent work using doses in the low clinically therapeutic range (low µM) showed amantadine to inhibit the N-methyl-D-aspartic acid (NMDA) receptor-mediated stimulation of acetylcholine release from rat striatum, most likely at the MK-801 site. Although amantadine does not possess anticholinergic activity in dogs at doses of 31.5 mg/kg, equivalent to an approximate human dose of 15.8 mg/kg (based on body surface area conversions), clinically, it exhibits anticholinergic-like side effects such as dry mouth, urinary retention, and constipation.

### Pharmacokinetics

SYMMETREL is well absorbed orally. Maximum plasma concentrations are directly related to dose for doses up to 200 mg/day. Doses above 200 mg/day may result in a greater than proportional increase in maximum plasma concentrations. It is primarily excreted unchanged in the urine by glomerular filtration and tubular secretion. Eight metabolites of amantadine have been identified in human urine. One metabolite, an N-acetylated compound, was quantified in human urine and accounted for 5-15% of the administered dose. Plasma acetylamantadine accounted for up to 80% of the concurrent amantadine plasma concentration in 5 of 12 healthy volunteers following the ingestion of a 200 mg dose of amantadine. Acetylamantadine was not detected in the plasma of the remaining seven volunteers. The contribution of this metabolite to efficacy or toxicity is not known.

There appears to be a relationship between plasma amantadine concentrations and toxicity. As concentration increases, toxicity seems to be more prevalent, however, absolute values of amantadine concentrations associated with adverse effects have not been fully defined.

Amantadine pharmacokinetics were determined in 24 normal adult male volunteers after the oral administration of a single amantadine hydrochloride 100 mg soft gel capsule. The mean ± SD maximum plasma concentration was 0.22 ± 0.03 µg/mL (range: 0.18 to 0.32 µg/mL). The time to peak concentration was 3.3 ± 1.5 hours (range: 1.5 to 8.0 hours). The apparent oral clearance was 0.28 ± 0.11 L/hr/kg (range: 0.14 to 0.62 L/hr/kg). The half-life was 17 ± 4 hours (range: 10 to 25 hours). Across other studies, amantadine plasma half-life has averaged 16 ± 6 hours (range: 9 to 31 hours) in 19 healthy volunteers.

After oral administration of a single dose of 100 mg amantadine syrup to five healthy volunteers, the mean ± SD maximum plasma concentration C<sub>max</sub> was 0.24 ± 0.04 µg/mL and ranged from

0.18 to 0.28 µg/mL. After 15 days of amantadine 100 mg b.i.d., the  $C_{max}$  was  $0.47 \pm 0.11$  µg/mL in four of the five volunteers. The administration of amantadine tablets as a 200 mg single dose to 6 healthy subjects resulted in a  $C_{max}$  of  $0.51 \pm 0.14$  µg/mL. Across studies, the time to  $C_{max}$  ( $T_{max}$ ) averaged about 2 to 4 hours.

Plasma amantadine clearance ranged from 0.2 to 0.3 L/hr/kg after the administration of 5 mg to 25 mg intravenous doses of amantadine to 15 healthy volunteers.

In six healthy volunteers, the ratio of amantadine renal clearance to apparent oral plasma clearance was  $0.79 \pm 0.17$  (mean  $\pm$  SD).

The volume of distribution determined after the intravenous administration of amantadine to 15 healthy subjects was 3 to 8 L/kg, suggesting tissue binding. Amantadine, after single oral 200 mg doses to 6 healthy young subjects and to 6 healthy elderly subjects has been found in nasal mucus at mean  $\pm$  SD concentrations of  $0.15 \pm 0.16$ ,  $0.28 \pm 0.26$ , and  $0.39 \pm 0.34$  µg/g at 1, 4, and 8 hours after dosing, respectively. These concentrations represented  $31 \pm 33\%$ ,  $59 \pm 61\%$ , and  $95 \pm 86\%$  of the corresponding plasma amantadine concentrations. Amantadine is approximately 67% bound to plasma proteins over a concentration range of 0.1 to 2.0 µg/mL. Following the administration of amantadine 100 mg as a single dose, the mean  $\pm$  SD red blood cell to plasma ratio ranged from  $2.7 \pm 0.5$  in 6 healthy subjects to  $1.4 \pm 0.2$  in 8 patients with renal insufficiency.

The apparent oral plasma clearance of amantadine is reduced and the plasma half-life and plasma concentrations are increased in healthy elderly individuals age 60 and older. After single dose administration of 25 to 75 mg to 7 healthy, elderly male volunteers, the apparent plasma clearance of amantadine was  $0.10 \pm 0.04$  L/hr/kg (range 0.06 to 0.17 L/hr/kg) and the half-life was  $29 \pm 7$  hours (range 20 to 41 hours). Whether these changes are due to decline in renal function or other age related factors is not known.

In a study of young healthy subjects (n=20), mean renal clearance of amantadine, normalized for body mass index, was significantly higher in males compared to females ( $p < 0.032$ ).

Compared with otherwise healthy adult individuals, the clearance of amantadine is significantly reduced in adult patients with renal insufficiency. The elimination half-life increases two to three fold or greater when creatinine clearance is less than 40 mL/min/1.73 m<sup>2</sup> and averages eight days in patients on chronic maintenance hemodialysis. Amantadine is removed in negligible amounts by hemodialysis.

The pH of the urine has been reported to influence the excretion rate of SYMMETREL. Since the excretion rate of SYMMETREL increases rapidly when the urine is acidic, the administration of urine acidifying drugs may increase the elimination of the drug from the body.

## **INDICATIONS AND USAGE**

SYMMETREL is indicated for the prophylaxis and treatment of signs and symptoms of infection caused by various strains of influenza A virus. SYMMETREL (Amantadine Hydrochloride) is also indicated in the treatment of parkinsonism and drug-induced extrapyramidal reactions.

**Influenza A Prophylaxis:** SYMMETREL is indicated for chemoprophylaxis against signs and symptoms of influenza A virus infection when early vaccination is not feasible or when the vaccine is contraindicated or not available. In the prophylaxis of influenza, early vaccination on an annual basis as recommended by the Centers for Disease Control's Immunization Practices Advisory Committee is the method of choice. Because SYMMETREL does not completely prevent the host immune response to influenza A infection, individuals who take this drug may still develop immune responses to natural disease or vaccination and may be protected when later exposed to antigenically related viruses. Following vaccination during an influenza A outbreak, SYMMETREL prophylaxis should be considered for the 2- to 4-week time period required to develop an antibody response.

**Influenza A Treatment:** SYMMETREL is also indicated in the treatment of uncomplicated respiratory tract illness caused by influenza A virus strains especially when administered early in the course of illness. There are no well-controlled clinical studies demonstrating that treatment with SYMMETREL will avoid the development of influenza A virus pneumonitis or other complications in high risk patients.

There is no clinical evidence indicating that SYMMETREL is effective in the prophylaxis or treatment of viral respiratory tract illnesses other than those caused by influenza A virus strains.

**Parkinson's Disease/Syndrome:** SYMMETREL is indicated in the treatment of idiopathic Parkinson's disease (Paralysis Agitans), postencephalitic parkinsonism, and symptomatic parkinsonism which may follow injury to the nervous system by carbon monoxide intoxication. It is indicated in those elderly patients believed to develop parkinsonism in association with cerebral arteriosclerosis. In the treatment of Parkinson's disease, SYMMETREL is less effective than levodopa, (-)-3-(3,4-dihydroxyphenyl)-L-alanine, and its efficacy in comparison with the anticholinergic antiparkinson drugs has not yet been established.

**Drug-Induced Extrapyrimal Reactions:** SYMMETREL is indicated in the treatment of drug-induced extrapyramidal reactions. Although anticholinergic-type side effects have been noted with SYMMETREL when used in patients with drug-induced extrapyramidal reactions, there is a lower incidence of these side effects than that observed with the anticholinergic antiparkinson drugs.

### **CONTRAINDICATIONS**

SYMMETREL is contraindicated in patients with known hypersensitivity to amantadine hydrochloride or to any of the other ingredients in SYMMETREL.

### **WARNINGS**

**Deaths:** Deaths have been reported from overdose with SYMMETREL. The lowest reported acute lethal dose was 1 gram. Acute toxicity may be attributable to the anticholinergic effects of amantadine. Drug overdose has resulted in cardiac, respiratory, renal or central nervous system toxicity. Cardiac dysfunction includes arrhythmia, tachycardia and hypertension (see OVERDOSAGE).

**Suicide Attempts:** Suicide attempts, some of which have been fatal, have been reported in patients treated with SYMMETREL, many of whom received short courses for influenza treatment or prophylaxis. The incidence of suicide attempts is not known and the pathophysiologic mechanism is not understood. Suicide attempts and suicidal ideation have been reported in patients with and without prior history of psychiatric illness. SYMMETREL can exacerbate mental problems in patients with a history of psychiatric disorders or substance abuse. Patients who attempt suicide may exhibit abnormal mental states which include disorientation, confusion, depression, personality changes, agitation, aggressive behavior, hallucinations, paranoia, other psychotic reactions, and somnolence or insomnia. Because of the possibility of serious adverse effects, caution should be observed when prescribing SYMMETREL to patients being treated with drugs having CNS effects, or for whom the potential risks outweigh the benefit of treatment.

**CNS Effects:** Patients with a history of epilepsy or other "seizures" should be observed closely for possible increased seizure activity.

Patients receiving SYMMETREL who note central nervous system effects or blurring of vision should be cautioned against driving or working in situations where alertness and adequate motor coordination are important.

**Other:** Patients with a history of congestive heart failure or peripheral edema should be followed closely as there are patients who developed congestive heart failure while receiving SYMMETREL.

Patients with Parkinson's disease improving on SYMMETREL should resume normal activities gradually and cautiously, consistent with other medical considerations, such as the presence of osteoporosis or phlebothrombosis.

Because SYMMETREL has anticholinergic effects and may cause mydriasis, it should not be given to patients with untreated angle closure glaucoma.

### **PRECAUTIONS**

SYMMETREL should not be discontinued abruptly in patients with Parkinson's disease since a few patients have experienced a parkinsonian crisis, i.e., a sudden marked clinical deterioration, when this medication was suddenly stopped. The dose of anticholinergic drugs or of SYMMETREL should be reduced if atropine-like effects appear when these drugs are used concurrently. Abrupt discontinuation may also precipitate delirium, agitation, delusions, hallucinations, paranoid reaction, stupor, anxiety, depression and slurred speech.

**Neuroleptic Malignant Syndrome (NMS):** Sporadic cases of possible Neuroleptic Malignant Syndrome (NMS) have been reported in association with dose reduction or withdrawal of SYMMETREL therapy. Therefore, patients should be observed carefully when the dosage of SYMMETREL is reduced abruptly or discontinued, especially if the patient is receiving neuroleptics.

NMS is an uncommon but life-threatening syndrome characterized by fever or hyperthermia; neurologic findings including muscle rigidity, involuntary movements, altered consciousness; mental status changes; other disturbances such as autonomic dysfunction, tachycardia, tachypnea, hyper- or hypotension; laboratory findings such as creatine phosphokinase elevation, leukocytosis, myoglobinuria, and increased serum myoglobin.

The early diagnosis of this condition is important for the appropriate management of these patients. Considering NMS as a possible diagnosis and ruling out other acute illnesses (e.g., pneumonia, systemic infection, etc.) is essential. This may be especially complex if the clinical presentation includes both serious medical illness and untreated or inadequately treated extrapyramidal signs and symptoms (EPS). Other important considerations in the differential diagnosis include central anticholinergic toxicity, heat stroke, drug fever and primary central nervous system (CNS) pathology.

The management of NMS should include: 1) intensive symptomatic treatment and medical monitoring, and 2) treatment of any concomitant serious medical problems for which specific treatments are available. Dopamine agonists, such as bromocriptine, and muscle relaxants, such as dantrolene are often used in the treatment of NMS, however, their effectiveness has not been demonstrated in controlled studies.

**Renal disease:** Because SYMMETREL is mainly excreted in the urine, it accumulates in the plasma and in the body when renal function declines. Thus, the dose of SYMMETREL should be reduced in patients with renal impairment and in individuals who are 65 years of age or older. Hemodialysis does not remove significant amounts of SYMMETREL; in patients with renal failure, a 4-hour hemodialysis removed 7 to 15 mg after a single 300 mg oral dose<sup>1</sup> (see DOSAGE AND ADMINISTRATION; Dosage for Impaired Renal Function).

**Liver disease:** Care should be exercised when administering SYMMETREL to patients with liver disease. Rare instances of reversible elevation of liver enzymes have been reported in patients receiving SYMMETREL, though a specific relationship between the drug and such changes has not been established.

**Other:** The dose of SYMMETREL may need careful adjustment in patients with congestive heart failure, peripheral edema, or orthostatic hypotension. Care should be exercised when administering SYMMETREL to patients with a history of recurrent eczematoid rash, or to patients with psychosis or severe psychoneurosis not controlled by chemotherapeutic agents.

Serious bacterial infections may begin with influenza-like symptoms or may coexist with or occur as complications during the course of influenza. SYMMETREL has not been shown to prevent such complications.

**Information for Patients:**

Patients should be advised of the following information:

Blurry vision and/or impaired mental acuity may occur.

Gradually increase physical activity as the symptoms of Parkinson's disease improve.

Avoid excessive alcohol usage, since it may increase the potential for CNS effects such as dizziness, confusion, lightheadedness and orthostatic hypotension.

Avoid getting up suddenly from a sitting or lying position. If dizziness or lightheadedness occurs, notify physician.

Notify physician if mood/mental changes, swelling of extremities, difficulty urinating and/or shortness of breath occur.

Do not take more medication than prescribed because of the risk of overdose. If there is no improvement in a few days, or if medication appears less effective after a few weeks, discuss with a physician.

Consult physician before discontinuing medication.

Seek medical attention immediately if it is suspected that an overdose of medication has been taken.

**Drug Interactions:** Careful observation is required when SYMMETREL is administered concurrently with central nervous system stimulants.

Agents with anticholinergic properties may potentiate the anticholinergic-like side effects of amantadine.

Coadministration of thioridazine has been reported to worsen the tremor in elderly patients with Parkinson's disease, however, it is not known if other phenothiazines produce a similar response. Coadministration of Dyazide (triamterene/hydrochlorothiazide) resulted in a higher plasma amantadine concentration in a 61-year-old man receiving SYMMETREL (Amantadine Hydrochloride) 100 mg TID for Parkinson's disease.<sup>2</sup> It is not known which of the components of Dyazide contributed to the observation or if related drugs produce a similar response.

Coadministration of trimethoprim-sulfamethoxazole may impair renal clearance of amantadine resulting in higher plasma concentrations.

Coadministration of quinine or quinidine with amantadine was shown to reduce the renal clearance of amantadine.

**Carcinogenesis and Mutagenesis:** Long-term *in vivo* animal studies designed to evaluate the carcinogenic potential of SYMMETREL have not been performed. In several *in vitro* assays for gene mutation, SYMMETREL did not increase the number of spontaneously observed mutations in four strains of *Salmonella typhimurium* (Ames Test) or in a mammalian cell line (Chinese Hamster Ovary cells) when incubations were performed either with or without a liver metabolic activation extract. Further, there was no evidence of chromosome damage observed in an *in vitro* test using freshly derived and stimulated human peripheral blood lymphocytes (with and without metabolic activation) or in an *in vivo* mouse bone marrow micronucleus test (140-550 mg/kg; estimated human equivalent doses of 11.7-45.8 mg/kg based on body surface area conversion).

**Impairment of Fertility:** In a three litter reproduction study in rats, SYMMETREL at a dose of 32 mg/kg/day (estimated human equivalent dose of 4.5 mg/kg/day, based on body surface area conversions) administered to both males and females slightly impaired fertility. There were no effects on fertility at a dose level of 10 mg/kg/day (estimated human equivalent dose of 1.4 mg/kg/day); intermediate doses were not tested.

Failed fertility has been reported during human *in vitro* fertilization (IVF) when the sperm donor ingested amantadine 2 weeks prior to, and during the IVF cycle.

**Pregnancy Category C:** SYMMETREL has been reported to be teratogenic in rats at 50 mg/kg/day and embryotoxic at 100 mg/kg/day (estimated human equivalent dose of 7.1 mg/kg/day and 14.2 mg/kg/day, respectively, based on body surface area conversion). A dose of 37 mg/kg/day (estimated human equivalent dose of 5.3 mg/kg/day) did not produce a teratogenic or embryotoxic effect in the rat. Embryotoxic and teratogenic effects were not seen in rabbits that received 32 mg/kg/day (estimated human equivalent dose of 9.6 mg/kg/day, based on body surface area conversion). There are no adequate and well-controlled studies in pregnant women. Human data regarding teratogenicity after maternal use of amantadine is scarce. Tetralogy of Fallot and tibial hemimelia (normal karyotype) occurred in an infant exposed to amantadine during the first trimester of pregnancy (100 mg P.O. for 7 days during the 6th and 7th week of gestation). Cardiovascular maldevelopment (single ventricle with pulmonary atresia) was associated with maternal exposure to amantadine (100 mg/d) administered during the first 2 weeks of pregnancy. SYMMETREL should be used during pregnancy only if the potential benefit justifies the potential risk to the embryo or fetus.

**Nursing Mothers:** SYMMETREL is excreted in human milk. Use is not recommended in nursing mothers.

**Pediatric Use:** The safety and efficacy of SYMMETREL in newborn infants and infants below the age of 1 year have not been established.

**Usage in the Elderly:** Because SYMMETREL is primarily excreted in the urine, it accumulates in the plasma and in the body when renal function declines. Thus, the dose of SYMMETREL should be reduced in patients with renal impairment and in individuals who are 65 years of age or older. The dose of SYMMETREL may need reduction in patients with congestive heart failure, peripheral edema, or orthostatic hypotension (see DOSAGE AND ADMINISTRATION).

## **ADVERSE REACTIONS**

The adverse reactions reported most frequently at the recommended dose of SYMMETREL (5-10%) are: nausea, dizziness (lightheadedness), and insomnia.

Less frequently (1-5%) reported adverse reactions are: depression, anxiety and irritability, hallucinations, confusion, anorexia, dry mouth, constipation, ataxia, livedo reticularis, peripheral edema, orthostatic hypotension, headache, somnolence, nervousness, dream abnormality, agitation, dry nose, diarrhea and fatigue.

Infrequently (0.1-1%) occurring adverse reactions are: congestive heart failure, psychosis, urinary retention, dyspnea, skin rash, vomiting, weakness, slurred speech, euphoria, thinking abnormality, amnesia, hyperkinesia, hypertension, decreased libido, and visual disturbance, including punctate subepithelial or other corneal opacity, corneal edema, decreased visual acuity, sensitivity to light, and optic nerve palsy.

Rare (less than 0.1%) occurring adverse reactions are: instances of convulsion, leukopenia, neutropenia, eczematoid dermatitis, oculogyric episodes, suicidal attempt, suicide, and suicidal ideation (see WARNINGS).

Other adverse reactions reported during postmarketing experience with SYMMETREL usage include:

**Nervous System/Psychiatric** - coma, stupor, delirium, hypokinesia, hypertonia, delusions, aggressive behavior, paranoid reaction, manic reaction, involuntary muscle contractions, gait abnormalities, paresthesia, EEG changes, and tremor. Abrupt discontinuation may also precipitate delirium, agitation, delusions, hallucinations, paranoid reaction, stupor, anxiety, depression and slurred speech;

**Cardiovascular** - cardiac arrest, arrhythmias including malignant arrhythmias, hypotension, and tachycardia;

**Respiratory** - acute respiratory failure, pulmonary edema, and tachypnea;

**Gastrointestinal** - dysphagia;

**Hematologic** - leukocytosis;

**Special Senses** - keratitis and mydriasis;

**Skin and Appendages** - pruritus and diaphoresis;

**Miscellaneous** - neuroleptic malignant syndrome (see WARNINGS), allergic reactions including anaphylactic reactions, edema, and fever;

**Laboratory Test** - elevated: CPK, BUN, serum creatinine, alkaline phosphatase, LDH, bilirubin, GGT, SGOT, and SGPT.

## **OVERDOSAGE**

Deaths have been reported from overdose with SYMMETREL. The lowest reported acute lethal dose was 1 gram. Because some patients have attempted suicide by overdosing with amantadine, prescriptions should be written for the smallest quantity consistent with good patient management.

Acute toxicity may be attributable to the anticholinergic effects of amantadine. Drug overdose has resulted in cardiac, respiratory, renal or central nervous system toxicity. Cardiac dysfunction includes arrhythmia, tachycardia and hypertension. Pulmonary edema and respiratory distress (including adult respiratory distress syndrome - ARDS) have been reported; renal dysfunction including increased BUN, decreased creatinine clearance and renal insufficiency can occur. Central nervous system effects that have been reported include insomnia, anxiety, agitation, aggressive behavior, hypertonia, hyperkinesia, ataxia, gait abnormality, tremor, confusion, disorientation, depersonalization, fear, delirium, hallucinations, psychotic reactions, lethargy, somnolence and coma. Seizures may be exacerbated in patients with prior history of seizure disorders. Hyperthermia has also been observed in cases where a drug overdose has occurred.

There is no specific antidote for an overdose of SYMMETREL. However, slowly administered intravenous physostigmine in 1 and 2 mg doses in an adult<sup>3</sup> at 1- to 2-hour intervals and 0.5 mg doses in a child<sup>4</sup> at 5- to 10-minute intervals up to a maximum of 2 mg/hour have been reported to be effective in the control of central nervous system toxicity caused by amantadine hydrochloride. For acute overdosing, general supportive measures should be employed along with immediate gastric lavage or induction of emesis. Fluids should be forced, and if necessary, given intravenously. Hemodialysis does not remove significant amounts of SYMMETREL; in patients with renal failure, a 4-hour hemodialysis removed 7 to 15 mg after a single 300 mg oral dose.<sup>1</sup> The pH of the urine has been reported to influence the excretion rate of SYMMETREL. Since the excretion rate of SYMMETREL increases rapidly when the urine is acidic, the administration of urine acidifying drugs may increase the elimination of the drug from the body. The blood pressure, pulse, respiration and temperature should be monitored. The patient should be observed for hyperactivity and convulsions; if required, sedation, and anticonvulsant therapy should be administered. The patient should be observed for the possible development of arrhythmias and hypotension; if required, appropriate antiarrhythmic and antihypotensive therapy should be given.



Electrocardiographic monitoring may be required after ingestion, since malignant tachyarrhythmias can appear after overdose.

Care should be exercised when administering adrenergic agents, such as isoproterenol, to patients with a SYMMETREL overdose, since the dopaminergic activity of SYMMETREL has been reported to induce malignant arrhythmias.

The blood electrolytes, urine pH and urinary output should be monitored. If there is no record of recent voiding, catheterization should be done.

## **DOSAGE AND ADMINISTRATION**

The dose of SYMMETREL may need reduction in patients with congestive heart failure, peripheral edema, orthostatic hypotension, or impaired renal function (see Dosage for Impaired Renal Function).

### **Dosage for Prophylaxis and Treatment of Uncomplicated Influenza A Virus Illness:**

**Adult:** The adult daily dosage of SYMMETREL (Amantadine Hydrochloride) is 200 mg; two 100 mg tablets (or four teaspoonfuls of syrup) as a single daily dose. The daily dosage may be split into one tablet of 100 mg (or two teaspoonfuls of syrup) twice a day. If central nervous system effects develop in once-a-day dosage, a split dosage schedule may reduce such complaints. In persons 65 years of age or older, the daily dosage of SYMMETREL is 100 mg.

A 100 mg daily dose has also been shown in experimental challenge studies to be effective as prophylaxis in healthy adults who are not at high risk for influenza-related complications. However, it has not been demonstrated that a 100 mg daily dose is as effective as a 200 mg daily dose for prophylaxis, nor has the 100 mg daily dose been studied in the treatment of acute influenza illness. In recent clinical trials, the incidence of central nervous system (CNS) side effects associated with the 100 mg daily dose was at or near the level of placebo. The 100 mg dose is recommended for persons who have demonstrated intolerance to 200 mg of SYMMETREL daily because of CNS or other toxicities.

**Pediatric Patients: 1 yr.-9 yrs. of age:** The total daily dose should be calculated on the basis of 2 to 4 mg/lb/day (4.4 to 8.8 mg/kg/day), but not to exceed 150 mg per day.

**9 yrs.-12 yrs. of age:** The total daily dose is 200 mg given as one tablet of 100 mg (or two teaspoonfuls of syrup) twice a day. The 100 mg daily dose has not been studied in this pediatric population. Therefore, there are no data which demonstrate that this dose is as effective as or is safer than the 200 mg daily dose in this patient population.

Prophylactic dosing should be started in anticipation of an influenza A outbreak and before or after contact with individuals with influenza A virus respiratory tract illness.

SYMMETREL should be continued daily for at least 10 days following a known exposure. If SYMMETREL is used chemoprophylactically in conjunction with inactivated influenza A virus vaccine until protective antibody responses develop, then it should be administered for 2 to 4 weeks after the vaccine has been given. When inactivated influenza A virus vaccine is unavailable or contraindicated, SYMMETREL should be administered for the duration of known influenza A in the community because of repeated and unknown exposure.

Treatment of influenza A virus illness should be started as soon as possible, preferably within 24 to 48 hours after onset of signs and symptoms, and should be continued for 24 to 48 hours after the disappearance of signs and symptoms.

### **Dosage for Parkinsonism:**

**Adult:** The usual dose of SYMMETREL is 100 mg twice a day when used alone. SYMMETREL has an onset of action usually within 48 hours.

The initial dose of SYMMETREL is 100 mg daily for patients with serious associated medical illnesses or who are receiving high doses of other antiparkinson drugs. After one to several weeks at 100 mg once daily, the dose may be increased to 100 mg twice daily, if necessary.

Occasionally, patients whose responses are not optimal with SYMMETREL at 200 mg daily may benefit from an increase up to 400 mg daily in divided doses. However, such patients should be supervised closely by their physicians.

Patients initially deriving benefit from SYMMETREL not uncommonly experience a fall-off of effectiveness after a few months. Benefit may be regained by increasing the dose to 300 mg daily. Alternatively, temporary discontinuation of SYMMETREL for several weeks, followed by reinitiation of the drug, may result in regaining benefit in some patients. A decision to use other antiparkinson drugs may be necessary.

**Dosage for Concomitant Therapy:** Some patients who do not respond to anticholinergic antiparkinson drugs may respond to SYMMETREL. When SYMMETREL or anticholinergic antiparkinson drugs are each used with marginal benefit, concomitant use may produce additional benefit.

When SYMMETREL and levodopa are initiated concurrently, the patient can exhibit rapid therapeutic benefits. SYMMETREL should be held constant at 100 mg daily or twice daily while the daily dose of levodopa is gradually increased to optimal benefit.

When SYMMETREL is added to optimal well-tolerated doses of levodopa, additional benefit may result, including smoothing out the fluctuations in improvement which sometimes occur in patients on levodopa alone. Patients who require a reduction in their usual dose of levodopa because of development of side effects may possibly regain lost benefit with the addition of SYMMETREL.

**Dosage for Drug-Induced Extrapyrimalidal Reactions:**

**Adult:** The usual dose of SYMMETREL is 100 mg twice a day. Occasionally, patients whose responses are not optimal with SYMMETREL at 200 mg daily may benefit from an increase up to 300 mg daily in divided doses.

**Dosage for Impaired Renal Function:**

Depending upon creatinine clearance, the following dosage adjustments are recommended:

CREATININE CLEARANCE (mL/min/1.73 m <sup>2</sup> )	SYMMETREL DOSAGE
30-50	200 mg 1st day and 100 mg each day thereafter
15-29	200 mg 1st day followed by 100 mg on alternate days
<15	200 mg every 7 days

The recommended dosage for patients on hemodialysis is 200 mg every 7 days.

**HOW SUPPLIED**

SYMMETREL is available in light orange, convex curved, triangular shaped 100 mg tablets with "SYMMETREL" debossed on one side and plain on the other side as follows:

Bottles of 100	NDC 63481-108-70
Bottles of 500	NDC 63481-108-85

As a clear, colorless syrup [each 5 mL (1 teaspoonful) contains 50 mg amantadine hydrochloride] in:

16 oz. (480 mL) bottles

NDC 63481-205-16

Store at 25°C (77°F), excursions permitted to 15°-30°C (59°-86°F). [See USP Controlled Room Temperature].

Dispense in a tight container as defined in the USP, with a child-resistant closure (as required).

#### REFERENCES

<sup>1</sup>V.W. Horadam, et. al., *Ann. Intern. Med.* 94:454, 1981.

<sup>2</sup>W.W. Wilson and A.H. Rajput, Amantadine-Dyazide Interaction, *Can. Med. Assoc. J.* 129:974-975, 1983.

<sup>3</sup>D.F. Casey, *N. Engl. J. Med.* 298:516, 1978.

<sup>4</sup>C.D. Berkowitz, *J. Pediatr.* 95:144, 1979.

Manufactured for:

**Endo Pharmaceuticals Inc.**

Chadds Ford, PA 19317

**Endo scripted logo**

Manufactured by:

DuPont Pharma

Wilmington, DE 19880

SYMMETREL® is a Registered Trademark of Endo Pharmaceuticals Inc.

Copyright © Endo Pharmaceuticals Inc. 2000

Printed in U.S.A.

6486-02/May, 2000